

9. Experimentelle Tests der Quantenelektrodynamik bei hohen Energien

9.1 Der Prozess $e^+e^- \rightarrow \mu^+\mu^-$


9.2 QED Streuprozesse

9.3 e^+e^- -Annihilation in Hadronen


9.4 Elastische Lepton-Nukleon-Streuung

9.5 Inelastische Lepton-Nukleon-Streuung

9.6 Das Quark-Parton Modell, Strukturfunktionen



Der totale Wirkungsquerschnitt für die Reaktion $e^+e^- \rightarrow \mu^+\mu^-$ als Funktion der Schwerpunktsenergie (aus Ref. [Halzen-Martin])



Die differentiellen Wirkungsquerschnitte für die Reaktionen $e^+e^- \rightarrow \mu^+\mu^-$ und $e^+e^- \rightarrow \tau^+\tau^-$ (aus Ref. [Schmüser])

e^+e^- Wirkungsquerschnitt






TABLE 6.1
 Leading Order Contributions to Representative QED Processes


	Feynman Diagrams		$ \overline{\mathcal{M}} ^2/2e^4$		
Møller scattering $e^-e^- \rightarrow e^-e^-$	Forward peak 	Backward peak 	Forward	Interference	Backward
(Crossing $s \leftrightarrow u$)			$\frac{s^2 + u^2}{t^2} + \frac{2s^2}{tu} + \frac{s^2 + t^2}{u^2}$		
			($u \leftrightarrow t$ symmetric)		
Bhabha scattering $e^-e^+ \rightarrow e^-e^+$	Forward 	"Time-like" 	Forward	Interference	Time-like
			$\frac{s^2 + u^2}{t^2} + \frac{2u^2}{ts} + \frac{u^2 + t^2}{s^2}$		
$e^- \mu^- \rightarrow e^- \mu^-$			$\frac{s^2 + u^2}{t^2}$		
(Crossing $s \leftrightarrow t$)			$\frac{u^2 + t^2}{s^2}$		






Differentielle Wirkungsquerschnitte für wichtige QED-Prozesse



Ratio R of (11.6) as a function of the total e^-e^+ center-of-mass energy. (The sharp peaks correspond to the production of narrow 1^- resonances just below or near the flavor thresholds.)




The proton form factors as a function of q^2 .





Comparison of the magnetic and electric form factors of neutron and proton. They are consistent with the scaling law (7.34). (a) Proton magnetic form factor; (b) proton electric form factor; (c) neutron magnetic form factor. (After Weber 1967.)


Das Proton für einen Atomphysiker




Das Proton für einen Kernphysiker




Das Proton für einen Teilchenphysiker






The $ep \rightarrow eX$ cross section as a function of the missing mass W . Data are from the Stanford Linear Accelerator. The elastic peak at $W = M$ has been reduced by a factor of 8.5.



The structure function νW_2 determined by electron-proton scattering as a function of Q^2 for $\omega = 4$. Data are from the Stanford Linear Accelerator.



The ratio $2xF_1/F_2$ measured in SLAC electron-nucleon scattering experiments. For spin- $\frac{1}{2}$ partons, with $g = 2$, a ratio of unity is expected in the limit of large q^2 – the Callan-Gross relation. (Data compiled from published SLAC data.)

Test der Callan-Gross-Relation



From where do we know the x-values?

The structure of the proton is investigated in *Deep Inelastic Scattering* experiments:


Highest energy machine: the HERA ep collider at DESY/Hamburg
Operated from 1990 - 2007

Scattering of 30 GeV electrons on 900 GeV protons:
→ Test of proton structure down to 10^{-18} m

HERA ep accelerator, 6.3 km circumference





Strukturfunktion $F_2(x, Q^2)$



- gemessen in Rutherford-Streuung an Eisen-Kernen für verschiedene x-Werte

heute: HERA - Speicherring @ DESY / Hamburg

$(e^- p \rightarrow e^- + X \dots)$
 $(27 \text{ GeV}) \times (820 \text{ GeV})$




Beispiel: ZEUS Experiment

$$Q^2 = 5300 \text{ GeV}^2$$

$$x = 0.11$$



1994





If the Proton is

then $F_2^{ep}(x)$ is



A quark





Three valence quarks




Three bound valence quarks



Three bound valence quarks + some slow debris, e.g., $g \rightarrow q\bar{q}$




Small x




A proton made up of valence quarks, gluons, and slow debris consisting of quark-antiquark pairs.


The structure function pictured corresponding to different compositions assumed for the proton.




The quark structure functions extracted from an analysis of deep inelastic scattering data. Figure (b) shows the total valence and sea quark contributions to the structure of the proton.




The ratio F_2^{en}/F_2^{ep} as a function of x , measured in deep inelastic scattering. Data are from the Stanford Linear Accelerator.



The difference $F_2^{ep} - F_2^{en}$ as a function of x , as measured in deep inelastic scattering. Data are from the Stanford Linear Accelerator.





Comparison of F_2^{vN} measured in neutrino-nucleon scattering in the Gargamelle heavy-liquid bubble chamber in a PS neutrino beam at CERN, with SLAC data on F_2^{eN} from electron-nucleon scattering, in the same region of q^2 . The data points are the neutrino results, and the curve is a fit through the electron data, multiplied by the factor $\frac{18}{5}$, which is the reciprocal of the mean squared charge of u - and d -quarks in the nucleon. This is a confirmation of the fractional charge assignments for the quarks. Note that the total area under the curve, measuring the total momentum fraction in the nucleon carried by quarks, is about 0.5. The remaining mass is ascribed to gluon constituents, which are the postulated carriers of the interquark color field



Momentum distributions of quarks (Q) and antiquarks (\bar{Q}) in the nucleon, at a value of q^2 of order 10 GeV^2 , obtained from results on neutrino and antineutrino scattering in experiments at CERN and Fermilab. The neutrino and antineutrino differential cross-sections measure the structure functions F_2 and F_3 in Eq. (7.50), and the difference and sum of these, through Eq. (7.56), give the quark and antiquark populations weighted by the momentum fraction x . The antiquarks (\bar{Q}) are concentrated at small x , the region of the so-called quark-antiquark "sea". The "valence" quarks of the static quark model ($Q - \bar{Q}$) are concentrated towards $x = 0.2$.

How do the x-values of the proton look like?



Parton density functions (pdf):

u- and d-quarks at large x-values

Glucos dominate at small x !!

Uncertainties in the pdfs,
 in particular on the gluon distribution
 at small x