


Physics at Hadron Colliders

Lecture 4



Search for Physics Beyond the Standard Model

- Supersymmetry
- Heavy particles decaying into di-leptons
- What if there are extra dimensions?

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Why do we think about extensions of the Standard Model ?

see lecture by E. Kiritsis

1. Gravity is not incorporated yet in the Standard Model
2. Many open questions in the Standard Model
 - Hierarchy problem: m_W (100 GeV) $\rightarrow m_{\text{Planck}}$ (10^{19} GeV)
 - Unification of couplings
 - Flavour / family problem
 -


All this calls for a **more fundamental theory** of which
the Standard Model is a low energy approximation \rightarrow **New Physics**

Candidate theories: Supersymmetry
Extra Dimensions
Technicolor
.....

All predict new physics at the TeV scale !!

Strong motivation for LHC mass reach ~ 3 TeV

The Search for Supersymmetry



K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Why do we like her so much?


1. Quadratically divergent quantum corrections to the Higgs boson mass are avoided

$$\Delta m_H = f(m_B^2 - m_f^2) \rightarrow m_{\text{SUSY}} \sim 1 \text{ TeV}$$


(Hierarchy or naturalness problem)

2. Unification of coupling constants of the three interactions seems possible

3. SUSY provides a candidate for dark matter,



The lightest SUSY particle (LSP)



4. A SUSY extension is a small perturbation, consistent with the electroweak precision data

the only problem:.....

No experimental evidence for SUSY so far ! (except that about half of the particles are already discovered)



Either SUSY does not exist

OR

m_{SUSY} large ($> 100 \text{ GeV}$) \rightarrow not accessible at present machines



LHC should say “final word” about (low energy) SUSY since theory predicts $m_{\text{SUSY}} \leq$ a few TeV

The Minimal Supersymmetric Standard Model (MSSM)

Symmetry between fermions (matter) and bosons (forces)

For each particle p with spin s , there exists a SUSY partner \tilde{p} with spin $\Delta s = 1/2$.

Ex. :	q ($s=1/2$)	\rightarrow	\tilde{q} ($s=0$)	squarks
	g ($s=1$)	\rightarrow	\tilde{g} ($s=1/2$)	gluino

Many new particles predicted !

Here : Minimal Supersymmetric extension of the Standard Model (MSSM)
which has minimal particle content



MSSM particle spectrum :

5 Higgs bosons : h, H, A, H \pm

quarks	\rightarrow	squarks	\tilde{q}
leptons	\rightarrow	sleptons	$\tilde{e}, \tilde{\mu}, \tilde{\nu}$, etc.
W^\pm	\rightarrow	winos	
H^\pm	\rightarrow	charged higgsino	$\left. \begin{array}{l} \rightarrow \chi^\pm_1, \chi^\pm_2 \\ \text{2 charginos} \end{array} \right\}$
γ	\rightarrow	photino	$\left. \begin{array}{l} \rightarrow \chi^0_{1,2,3,4} \\ \text{4 neutralinos} \end{array} \right\}$
Z	\rightarrow	zino	
h, H	\rightarrow	neutral higgsino	
g	\rightarrow	gluino	\tilde{g}

SUSY phenomenology

There is a multiplicative quantum number:

$$\text{R-parity} \quad R_p = \begin{cases} +1 & \text{Standard Model particles} \\ -1 & \text{SUSY particles} \end{cases}$$

which is **conserved** in most popular models (considered here).

Consequences:

- SUSY particles are **produced in pairs**
- **Lightest Supersymmetric Particle (LSP) is stable.**
In most models LSP is also **weakly interacting**:


$$\boxed{\text{LSP} = \chi^0_1}$$

- LSP is a good candidate for cold **dark matter**
- LSP behaves like a v → it escapes detection
- E_T^{miss} (typical SUSY signature)

The **masses of the SUSY particles** are not predicted;
Theory has many additional new parameters (on which the masses depend)

However, charginos/neutralinos are usually lighter than squarks/sleptons/gluinos.

Present mass limits : m (sleptons, charginos) > 90-103 GeV LEP II
m (squarks, gluinos) > ~ 250 GeV Tevatron Run 1
m (LSP, lightest neutralino) > ~ 45 GeV LEP II




K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Search for Supersymmetry at the LHC

- If **SUSY** exists at the electroweak scale, a discovery at the LHC should be easy
- Squarks and Gluinos** are strongly produced

They decay through cascades to the lightest SUSY particle (LSP)




- Step: Look for **deviations from the Standard Model**
Example: Multijet + E_T^{miss} signature
- Step: Establish the **SUSY mass scale** use inclusive variables, e.g. effective mass distribution
- Step: Determine **model parameters** (difficult)
Strategy: select particular decay chains and use kinematics to determine mass combinations

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Squarks and Gluinos

- Strongly produced, cross sections comparable to QCD cross sections at the same mass scale
- If R-parity conserved, cascade decays produce distinctive events: multiple jets, leptons, and E_T^{miss}
- Typical selection: $N_{\text{jet}} > 4$, $E_T > 100, 50, 50, 50$ GeV, $E_T^{\text{miss}} > 100$ GeV
- Define: $M_{\text{eff}} = E_T^{\text{miss}} + P_T^1 + P_T^2 + P_T^3 + P_T^4$ (effective mass)



LHC reach for Squark- and Gluino masses:

$$\begin{aligned} 1 \text{ fb}^{-1} &\Rightarrow M \sim 1500 \text{ GeV} \\ 10 \text{ fb}^{-1} &\Rightarrow M \sim 1900 \text{ GeV} \\ 100 \text{ fb}^{-1} &\Rightarrow M \sim 2500 \text{ GeV} \end{aligned}$$


TeV-scale SUSY can be found quickly !

K. Jakobs, Universität Freiburg


CERN Summer Student Lectures, Aug. 2005

LHC reach in the m_0 - $m_{1/2}$ mSUGRA plane:

Multijet + E_T^{miss} signature



SUSY cascade decays give also rise to many other inclusive signatures: leptons, b-jets, τ 's





Expect multiple signatures for TeV-scale SUSY

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Determination of model parameters

- Invisible LSP \Rightarrow no mass peaks, but kinematic endpoints
 \Rightarrow mass combinations
- Simplest case: $\chi^0_2 \rightarrow \chi^0_1 \ell^+ \ell^-$ endpoint: $M_{\ell\ell} = M(\chi^0_2) - M(\chi^0_1)$
 \sim
 (significant mode if no $\chi^0_2 \rightarrow \chi^0_1 Z, \chi^0_1 h, \ell^+ \ell^-$ decays)
- Require: 2 isolated leptons, multiple jets, and large E_T^{miss}





Modes can be distinguished
using shape of $\ell\ell$ -spectrum

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

h $\rightarrow bb:$



important if $\chi^0_2 \rightarrow \chi^0_1 h$ is open;
bb peak can be reconstructed in
many cases

Could be a Higgs discovery mode !

SM background can be reduced
by applying a cut on E_T^{miss}

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Strategy in SUSY Searches at the LHC:

- Search for multijet + E_T^{miss} excess
- If found, select SUSY sample (simple cuts)
- Look for special features (γ 's, long lived sleptons)
- Look for ℓ^\pm , $\ell^+ \ell^-$, $\ell^\pm \ell^\pm$, b-jets, τ 's
- End point analyses, global fit



K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005


Status of SUSY Searches at the Tevatron - a few examples-

(1) Search for Charginos and Neutralinos

- These particles can decay via W^* / Z^* or sleptons into **Leptons + P_T^{miss}**

Example: associated $\tilde{\chi}_0^0 \tilde{\chi}_1^\pm$ production

$$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow l^\pm l^\mp l^\pm \tilde{\chi}_1^0 \tilde{\chi}_1^0 X$$




- Small background from Standard Model processes (**striking signature**), but also **a small signal cross section**
- Leptonic branching ratios are enhanced if slepton masses are close to gaugino masses



Analysis: Search for 5 ($\ell\ell\ell$) + like-sign di-muon final states with missing transverse momentum

Selection	background expected	# obs. events
ee+l	0.21±0.12	0
eμ+l	0.31±0.13	0
μμ+l	1.75±0.57	2
μ [±] μ [±]	0.64±0.38	1
eτ+l	0.58±0.14	0
μτ+l	0.36±0.13	1
SUM	3.85±0.75	4

mSUGRA interpretation




For specific scenarios: sensitivity / limits above LEP limits

K. Jakobs, Universität Freiburg


CERN Summer Student Lectures, Aug. 2005

(2) Search for Squarks and Gluinos

- Copiously produced (QCD production)



- Cascade decays into multi-jet + P_T^{miss} final states
(2,3 or 4 jets, depending on squark / gluino mass relations)




Limits beyond LEP mSUGRA reach for $m(\text{squark}) \sim m(\text{gluino})$

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

The Reach for SUSY at the Tevatron

SUSY:
Extend reach for SUSY
particle masses
(for 10 fb⁻¹)



K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005



Can LHC probe extra dimensions ?


- Much recent theoretical interest in models with extra dimensions
(Explain the weakness of gravity (or hierarchy problem) by extra dimensions)
- New physics can appear at the TeV-mass scale,
i.e. accessible at the LHC

Example: Search for direct Graviton production

$$gg \rightarrow gG, qg \rightarrow qG, q\bar{q} \rightarrow Gg$$

$$q\bar{q} \rightarrow G\gamma$$

\Rightarrow Jets or Photons with E_T^{miss}




K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Search for escaping gravitons:

Jet + E_T^{miss} search:



$$G_N^{-1} = 8\pi R^\delta M_D^{2+\delta}$$

δ : # extra dimensions
 M_D = scale of gravitation
 R = radius (extension)

M_D^{max}	=	9.1, 7.0, 6.0 TeV
for		
δ	=	2, 3, 4
Extension: $10^{-5}, 10^{-10}, 10^{-12} \text{ m}$		

Main backgrounds:
jet+Z($\rightarrow vv$), jet+W \rightarrow jet+(e, μ , τ)v

„LHC experiments are also sensitive to this field of physics“ \rightarrow robust detectors

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

More ideas?

1. What about heavy new resonances decaying into lepton pairs

examples: W' and Z' or Graviton resonances (extra dimensions)

use again leptonic decay mode to search for them: $W' \rightarrow \ell \nu$
 $Z' \rightarrow \ell \ell$

Increased sensitivity in the Tevatron Run II

2. What about Leptoquarks ?

Particles that decay into leptons and quarks
(violate lepton and baryon number; appear in Grand Unified theories)



here: search for low mass Leptoquarks (TeV scale)

Fermilab Search for New Resonances in High Mass Di-leptons

- Neutral Gauge Boson Z'
◆ SM Coupling assumed

- Randall-Sundrum narrow Graviton resonances decaying to di-lepton

appear in Extra Dim. Scenarios




Main background from Drell-Yan pairs


Search for New Resonances in High Mass Di-leptons



Di-electron Invariant Mass



Di-muon Invariant Mass



Data are consistent with background from SM processes. No excess observed.

Z' mass limits (SM couplings)

95% C.L.

CDF :

DØ:

ee

μμ

ee+μμ

ττ

GeV/c²

750

735

815

394

780

680

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005




Search for W' → eν

- W': additional charged heavy vector boson
- appears in theories based on the extension of the gauge group
- e.g. Left-right symmetric models: $SU(2)_R \times W_R$
- assume: the neutrino from W' decay is light and stable.
- signature:

high p_T electron + high P_T^{miss}

→ peak in transverse mass distribution




K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005



Search for $W' \rightarrow e\nu$



Data:

consistent with one well known W
+ background



Limit: $M(W') > 842 \text{ GeV}/c^2$


(assuming Standard Model couplings)

K. Jakobs, Universität Freiburg


CERN Summer Student Lectures, Aug. 2005

Search for Scalar Leptoquarks (LQ)

- Production:
pair production via QCD processes
($q\bar{q}$ and gg fusion)
- Decays: into a lepton and a quark



$\beta =$ LQ branching fraction to charged lepton and quark
 $N =$ generation index
Leptoquarks of 1., 2., and 3. generation



Experimental Signatures:

- two high p_T isolated leptons + jets .OR.
- one isolated lepton + $P_T^{\text{miss}} + \text{jets}$.OR.
- $P_T^{\text{miss}} + \text{jets}$

K. Jakobs, Universität Freiburg


CERN Summer Student Lectures, Aug. 2005




1st, 2nd and 3rd generation Leptoquarks



channels: eejj, ev jj



channels: μμjj, evjj, vvjj



95% C.L. Mass Limits	1. Generation LQ	2. Generation LQ	3. Generation LQ
CDF (Run II)	235 GeV/c ²	224 GeV/c ²	129 GeV/c ²
D0 (Run I + II)	256 GeV/c ²	200 GeV/c ² (Run I)	


K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

LHC reach for other BSM Physics (a few examples for 30 and 100 fb⁻¹)

	30 fb ⁻¹	100 fb ⁻¹
Excited Quarks $Q^* \rightarrow q \gamma$	$M(q^*) \sim 3.5 \text{ TeV}$	$M(q^*) \sim 6 \text{ TeV}$
Leptoquarks	$M(LQ) \sim 1 \text{ TeV}$	$M(LQ) \sim 1.5 \text{ TeV}$
$Z' \rightarrow l\bar{l}, jj$ $W' \rightarrow l\nu$	$M(Z') \sim 3 \text{ TeV}$ $M(W') \sim 4 \text{ TeV}$	$M(Z') \sim 5 \text{ TeV}$ $M(W') \sim 6 \text{ TeV}$
Compositeness (from Di-jet)	$\Lambda \sim 25 \text{ TeV}$	$\Lambda \sim 40 \text{ TeV}$

$$\int \mathcal{L} dt = 100 \text{ fb}^{-1}$$



K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

Conclusions

1. Experiments at Hadron Colliders have a huge discovery potential
 - **SM Higgs:** full mass range, already at low luminosity
Vector boson fusion channels improve the sensitivity significantly
 - **MSSM Higgs:** parameter space covered
 - **SUSY:** discovery of TeV-scale SUSY should be easy,
determination of model parameters is more difficult
 - **Exotics:** experiments seem robust enough to cope with new scenarios
2. Experiments have also a great potential for precision measurements
 - m_W to ~ 15 MeV
 - m_{top} to ~ 1 GeV
 - $\Delta m_H / m_H$ to 0.1% (100 - 600 GeV)
 - + gauge couplings and measurements in the top sector

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

LHC : most difficult and ambitious high-energy physics project ever realized
(human and financial resources, technical challenges, complexity,)

It has a crucial role in physics: can say the final word about

- SM Higgs mechanism
- low-energy SUSY and other TeV-scale predictions



It will most likely modify our understanding of Nature

There are very exciting times ahead of us !!

We hope that many of you will join us in the discovery enterprise

- In case you have any questions:
please do not hesitate to contact me: karl.jakobs@uni-freiburg.de
- Transparencies will be made available as .pdf files on the web
(official summer school pages)

K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005

End of lectures



K. Jakobs, Universität Freiburg

CERN Summer Student Lectures, Aug. 2005