



# *Entdeckung eines Higgs-artigen Teilchens am LHC*



Karl Jakobs  
Physikalisches Institut  
Universität Freiburg

# *Entdeckung eines Higgs-artigen Teilchens am LHC*



*The ATLAS and CMS Collaborations*




**4<sup>th</sup> July 2012**





From the editorial:

*The top Breakthrough of the Year – the discovery of the Higgs boson – was an unusually easy choice, representing both a triumph of the human intellect and the culmination of decades of work by many thousands of physicists and engineers*



# a New Particle

Submission to PLB on 31<sup>st</sup> July 2012



Contents lists available at SciVerse ScienceDirect

Physics Letters B

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)



Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC <sup>☆</sup>

ATLAS Collaboration \*

This paper is dedicated to the memory of our ATLAS colleagues who did not live to see the full impact and significance of their contributions to the experiment.



Contents lists available at SciVerse ScienceDirect

Physics Letters B

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)



Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC <sup>☆</sup>



CMS Collaboration \*

CERN, Switzerland

This paper is dedicated to the memory of our colleagues who worked on CMS but have since passed away. In recognition of their many contributions to the achievement of this observation.

Decay observed into particles with same spin and electric charge sum = 0  
→ a new neutral boson has been discovered

# The Standard Model of Particle Physics




$$m_W \approx 80.4 \text{ GeV}$$

$$m_Z \approx 91.2 \text{ GeV}$$


- (i) Constituents of matter: quarks and leptons (spin-½ fermions)
- (ii) Four fundamental forces: described by quantum field theories (except gravitation)
  - massless spin-1 gauge bosons
- (iii) The Higgs field:
  - scalar field, spin-0 Higgs boson

# The Brout-Englert-Higgs Mechanism



F. Englert and R. Brout. Phys. Rev. Lett. 13 (1964) 321;  
P.W. Higgs, Phys. Lett. 12 (1964) 132, Phys. Rev. Lett. 13 (1964) 508;  
G.S. Guralnik, C.R. Hagen, and T.W.B. Kibble. Phys. Rev. Lett. 13 (1964) 585.

# The Brout-Englert-Higgs Mechanism




Complex scalar (spin 0) field  $\phi$  with potential:

$$V(\phi) = \mu^2(\phi^* \phi) + \lambda(\phi^* \phi)^2$$

For  $\lambda > 0$ ,  $\mu^2 < 0$ :  
“Spontaneous Symmetry Breaking”

- Omnipresent Higgs field: vacuum expectation value  $v \approx 246$  GeV
- **Higgs Boson** (mass not predicted, except  $m_H < \sim 1000$  GeV)
- Particles acquire mass through couplings to the Higgs field

# The Brout-Englert-Higgs Mechanism





Complex scalar (spin 0) field  $\phi$  with potential:

$$V(\phi) = \mu^2(\phi^* \phi) + \lambda(\phi^* \phi)^2$$

For  $\lambda > 0$ ,  $\mu^2 < 0$ :

“Spontaneous Symmetry Breaking”




- Couplings proportional to mass
- Higgs boson decays preferentially into the heaviest accessible particles

The Higgs field solves two fundamental problems:

(i) Masses of the vector bosons W and Z and fermions


(ii) Divergences in the theory (scattering of W bosons)  
("Ultraviolet regulator")



$$-iM(W^+W^- \rightarrow W^+W^-) \sim \frac{s}{M_W^2} \quad \text{for} \quad s \rightarrow \infty \quad (\text{no Higgs boson})$$

$$-iM(W^+W^- \rightarrow W^+W^-) \sim m_H^2 \quad \text{for} \quad s \rightarrow \infty \quad (\text{with Higgs boson})$$

# *The Large Hadron Collider*



CMS



LHCb




ALICE



ATLAS

# The ATLAS experiment




Diameter	25 m
Barrel toroid length	26 m
End-cap end-wall chamber span	46 m
Overall weight	7000 Tons


ATLAS Germany:  
BMBF-Forschungsschwerpunkt ATLAS



- HU Berlin, Bonn, DESY, Dortmund, Dresden, Freiburg, Giessen, Göttingen, Heidelberg, Mainz, LMU München, MPI München, Siegen, Würzburg, Wuppertal
- ~ 420 scientists (~200 students)



# The CMS experiment



Total weight 12500 t  
Overall diameter 15 m  
Overall length 21.6 m

Drift Tube Chambers (DT)


Resistive P Chambers

CMS Germany:  
BMBF-Forschungsschwerpunkt CMS


Aachen (Ib, IIIa, IIIb), DESY,  
Hamburg, Karlsruhe

~ 200 scientists (~90 students)





# Data taking in 2011/ 2012



Some results with the full dataset

Until end 2012:


$> 10^{15}$  pp collisions

$\sim 10^{10}$  pp collisions recorded



$25 \cdot 10^6 Z \rightarrow \mu\mu$  decays produced

- Excellent LHC performance  
Peak luminosities  $> 7 \cdot 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$  (world record, 2012)
- Excellent performance of the experiments:
  - Data recording efficiency  $\sim 93.5\%$
  - Working detector channels  $> 99\%$
  - Speed of data analysis


# The Standard Model at the LHC




# Higgs Boson Production




Gluon fusion



Vector boson  
fusion




WH/ZH  
associated  
production



$t\bar{t}H$   
associated  
production

\*) LHC Higgs cross-section working group  
Large theory effort


# Higgs Boson Decays



Important channels at hadron colliders:

- $H \rightarrow WW \rightarrow \ell^+ \ell^- \nu \bar{\nu}$
- $H \rightarrow \gamma\gamma$
- $H \rightarrow ZZ \rightarrow \ell^+ \ell^- \ell^+ \ell^-$


# Discovery of a Higgs-like particle



Expected number of decays in data:  
 $m_H = 125 \text{ GeV}$

- ~ 950  $H \rightarrow \gamma\gamma$
- ~ 60  $H \rightarrow ZZ \rightarrow 4\ell$
- ~ 9000  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$


# Search for the $H \rightarrow \gamma\gamma$ decay





- 2 photons (isolated) with large transverse momenta
- Mass of the Higgs boson can be reconstructed  $m_{\gamma\gamma}$

Both experiments have a good mass resolution  
 ATLAS:  $\sim 1.7 \text{ GeV}/c^2$  for  $m_H \sim 120 \text{ GeV}/c^2$

- Challenges:
  - signal-to-background ratio  
 (small, but smooth irreducible  $\gamma\gamma$  background)





- reducible backgrounds from  $\gamma j$  and  $jj$   
 (several orders of magnitude larger than irreducible one)





# Result of the ATLAS search for $H \rightarrow \gamma\gamma$





- $p_0$  value for consistency of data with background-only:  $\sim 10^{-13}$  ( $7.4\sigma$ )

More details: "Eingeladener Vortrag" by Kerstin Tackmann (Thursday)  
(Hertha-Sponer Preisträgerin)



# Result of the ATLAS search for $H \rightarrow \gamma\gamma$



Mass:






$$m_H = 126.8 \pm 0.2 \text{ (stat)} \pm 0.7 \text{ (syst)} \text{ GeV}$$

Signal strength:




$$\mu := \sigma / \sigma_{\text{SM}} = 1.65 \pm 0.24 \text{ (stat)} {}^{+0.17}_{-0.13} \text{ (syst)} {}^{+0.18}_{-0.13} \text{ (theo)}$$





## Separation of different production processes for $H \rightarrow \gamma\gamma$




# Result of the CMS search for $H \rightarrow \gamma\gamma$



# Search for the $H \rightarrow ZZ^{(*)} \rightarrow \ell^+\ell^- \ell^+\ell^-$ decay



- The “golden mode”  
4 leptons (isolated) with large transverse momenta
- Mass of the Higgs boson can be reconstructed  $m_{4\ell}$   
Both experiments have a good mass resolution  
ATLAS: ~2.5 GeV/c<sup>2</sup> (4e) for  $m_H \sim 130$  GeV/c<sup>2</sup>  
~2.0 GeV/c<sup>2</sup> (4μ) for  $m_H \sim 130$  GeV/c<sup>2</sup>




- Low signal rate, but also low background
  - Mainly from ZZ continuum

- In addition from tt and Zbb events:  
 $tt \rightarrow Wb Wb \rightarrow \ell\nu c\bar{\ell}\nu \ell\nu c\bar{\ell}\nu$   
 $Z bb \rightarrow \ell\ell c\bar{\ell}\nu c\bar{\ell}\nu$

# Candidate event for a $H \rightarrow ZZ \rightarrow e^+e^- \mu^+\mu^-$ decay


**ATLAS**  
EXPERIMENT  
<http://atlas.ch>




# 4 $\ell$ invariant mass spectra



Full dataset



Full dataset




- $p_0$ -values in both experiments  
 $\sim 10^{-11}$  ( $> 6\sigma$ )

Signal strengths:


ATLAS:  $\mu = 1.7 \pm 0.5$

CMS:  $\mu = 0.91^{+0.30}_{-0.24}$


# Time evolution of the $H \rightarrow ZZ \rightarrow 4l$ signal




# Time evolution of the $H \rightarrow ZZ \rightarrow 4\ell$ signal




# Time evolution of the $H \rightarrow ZZ \rightarrow 4l$ signal



# Separation of different production processes for $H \rightarrow ZZ \rightarrow 4\ell$



# Results on the search for $H \rightarrow WW \rightarrow l\nu l\nu$ decays




Signal strength:

ATLAS:  $\mu = 1.4 \pm 0.3$


CMS:  $\mu = 0.76 \pm 0.21$

Full dataset




Clear excess in both experiments

# Determination of mass, compatibility of channels



$$m_H = 125.2 \pm 0.3 \text{ (stat)} \pm 0.6 \text{ (syst)} \text{ GeV}$$



$$m_H = 125.8 \pm 0.4 \text{ (stat)} \pm 0.4 \text{ (syst)} \text{ GeV}$$


Updated mass values (full dataset, preliminary):

$$m_H (\gamma\gamma) = 126.8 \pm 0.2 \text{ (stat)} \pm 0.7 \text{ (syst)} \text{ GeV}$$

$$m_H (4\ell) = 124.3^{+0.6}_{-0.5} \text{ (stat)}^{+0.5}_{-0.4} \text{ (syst)} \text{ GeV}$$

$$m_H (4\ell) = 125.8 \pm 0.5 \text{ (stat)} \pm 0.2 \text{ (syst)} \text{ GeV}$$

# *Couplings to quarks and leptons ?*




Search for  $H \rightarrow \tau\tau$  and  $H \rightarrow bb$  decays

# Why is the search in these decay modes so challenging?

- The  $\tau$  lepton is the heaviest lepton


$$m_\tau = 1.78 \text{ GeV } /c^2, \text{ lifetime } 2.9 \cdot 10^{-13} \text{ s}$$


Decays into hadrons  $\tau \rightarrow \text{hadrons } \nu_\tau$  (65%)  
 $\tau \rightarrow e\nu_e\nu_\tau, \mu\nu_\mu\nu_\tau$  (35%)



- Challenge: distinguish hadronic  $\tau$  decays from hadronic jet activity

- Neutrinos in the final state, poor mass resolution





- Analysis is split into three sub-channels:
  - $H \rightarrow \tau\tau \rightarrow \ell\nu\nu \quad \ell\nu\nu$
  - $H \rightarrow \tau\tau \rightarrow \ell\nu\nu \quad \text{had } \nu$
  - $H \rightarrow \tau\tau \rightarrow \text{had } \nu \quad \text{had } \nu$
- Data set:  $13 - 17 \text{ fb}^{-1}$





Signal strength (all sub-channels):

ATLAS:  $\mu = 0.7 \pm 0.7$

CMS:  $\mu = 0.7 \pm 0.5$



More details: “Eingeladener Vortrag” by Stanley Lai (Thursday)

## Results on $H \rightarrow bb$ searches



- Small excess is showing up around 125 GeV
  - However, the significance is still low !

# Results on $H \rightarrow b\bar{b}$ from ATLAS



Signal strength:

ATLAS:  $\mu = -0.4 \pm 1.0$

CMS:  $\mu = 1.3^{+0.7}_{-0.6}$




# *Is the new particle the Higgs Boson ?*

- Production rates ?

Couplings to bosons and fermions

- Spin,  $J^P$  quantum number





More details: “*Eingeladener Vortrag*” by Johannes Elmsheuser (Thursday)  
“*Hauptvortrag*” by Thomas Müller (Friday)



# Signal strength in individual decay modes

-including new data up to Sept. 2012-

updates expected  
very soon




- Data are consistent with the hypothesis of a Standard Model Higgs boson !
- Experimental uncertainties are still too large to get excited about “high”  $\gamma\gamma$  and “low” fermionic ( $\tau\tau$  and  $bb$ ) signal strengths !

# Test of coupling strengths

updates expected  
very soon




## Couplings to W and Z bosons



$\lambda_{WZ}$  = ratio of W/Z coupling strength  
(normalized to ratio in the SM)

$\kappa_Z$  = scale factor for Z coupling strength


## W,Z versus fermion couplings



$\kappa_V$  = common scale factor for all fermion  
couplings (t, b,  $\tau$ , ....)

$\kappa_V$  = common scale factor for W,Z couplings


# Spin and CP





*Wolfgang Pauli und Niels Bohr bei der  
wissenschaftlichen Untersuchung  
der Kreiselbewegung  
(1952, anlässlich der Eröffnung des Instituts  
für Theoretische Physik in Lund / Schweden)*

- If Standard Model Higgs boson:  $J^P = 0^+$   
→ strategy is to falsify other hypotheses  
( $0^-$ ,  $1^-$ ,  $1^+$ ,  $2^-$ ,  $2^+$ )  
  
Spin 1: dis-favoured by observed  
 $H \rightarrow \gamma\gamma$  decays, Landau-Yang theorem  
  
Spin 2: consider graviton-like tensor,  
equivalent to a Kaluza-Klein graviton
- Angular distributions of final state  
particles show sensitivity to spin

# Spin studies using $H \rightarrow \gamma\gamma$ events



Decay angle in the Higgs boson rest frame  
(Collins-Soper frame)



$\cos \theta^*$  distribution in signal region, after background subtraction



Likelihood hypothesis test of spin-0 versus spin 2:



Data favour spin 0,

exclude  $J^P = 2^+$  w.r.t  $J^P = 0^+$  with

ATLAS:  $1.4\sigma$  (p value = 0.12)

# Spin studies using $H \rightarrow ZZ^{(*)} \rightarrow 4l$ events

- Sensitive variables:
  - Masses of the two Z bosons
  - Production angle  $\theta^*$
  - Four decay angles  $\Phi_1, \Phi, \theta_1$  and  $\theta_2$



Updated with full dataset  
 Likelihood hypothesis test of

$J^P = 0^+$  vs.  $J^P = 0^-$  ATLAS:  $2.8\sigma$  (2.7 exp)  
 CMS:  $3.3\sigma$  (2.6 exp)

$J^P = 0^+$  vs.  $J^P = 2^+_m$  ATLAS:  $1.2\sigma$  (1.5 exp)  
 CMS:  $2.7\sigma$  (1.8 exp)

Data favour  $J^P = 0^+$  w.r.t. other  $J^P$  configurations (including  $0^-, 1^-, 1^+, 2^+$ )



# Conclusions

- With the operation of the LHC at high energies, particle physics has entered a new era
- Performance of the LHC and the experiments is superb
- A milestone discovery made in July 2012

Strong evidence that the new particle is the long-sought Higgs boson

- Clear signals in bosonic decays
  - Evidence for fermionic decays is building up, but still weak....
  - Tensions in some signal/coupling strength (?)
  - First evidence for spin 0
- 
- 

- Exciting times ahead of us:
  - Study of the Higgs-like boson itself
  - Search for Physics Beyond the Standard Model

