


# *Experimental Status and Recent Progress on Electroweak Symmetry Breaking*



Karl Jakobs, University of Freiburg

-Results from the ATLAS and CMS Collaborations-





## *"Summary of Results from LHC Run 1"*



- Present status on:
  - Bosonic decay modes  
 $H \rightarrow \gamma\gamma$ ,  $H \rightarrow ZZ^*$ ,  $H \rightarrow WW^*$
  - Decays into fermions
  - Search for rare decays
- Profile of the new particle  
(mass, Spin-CP, couplings)
- Additional Higgs bosons?
- Prospects for LHC Run 2



Steve Myers PLHC 2012:

"The first two years of LHC operation have produced sensational performance: well beyond our wildest expectations. The combination of the performance of the LHC machine, the detectors and the GRID have proven to be a terrific success story in particle physics."

# Performance of the LHC and of the experiments



- Excellent LHC performance in 2011 and 2012
- Peak luminosities  $> 7 \cdot 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
- High level of pileup: mean of  $\sim 20$  interactions / beam crossing in 2012
- Excellent performance of the ATLAS and CMS experiments:  
(Data recording efficiency:  $\sim 93.5\%$ , working detector channels  $> 97\%$  for most sub-detectors, high data quality, speed of the data analysis)




## Standard Model Production Cross Section Measurements

Status: March 2015

ATLAS Preliminary


Run 1  $\sqrt{s} = 7, 8 \text{ TeV}$







## Standard Model Production Cross Section Measurements

Status: March 2015




“Stairway to Heaven”


# Higgs Boson Production




Gluon fusion



Vector boson  
fusion



WH/ZH  
associated  
production




$t\bar{t}$  associated  
production

\*) LHC Higgs cross-section working group  
Large theory effort

Meanwhile the NNNLO =  $N^3LO$  calculation for the gluon-fusion process exists;  
B. Anastasiou et al. (2015) → LHC = Long and Hard Calculations

# Higgs Boson Decays



Useful decays at a hadron collider:

- Final states with **leptons** via WW and ZZ decays
- **$\gamma\gamma$  final states** (despite small branching ratio)
- $\tau\tau$  final states (more difficult)
- In addition:  $H \rightarrow bb$  decays via associated lepton signatures (VBF, VH or ttH production)

SM predictions ( $m_H = 125.5$  GeV):


$$\begin{aligned} BR(H \rightarrow WW) &= 22.3\% \\ BR(H \rightarrow ZZ) &= 2.8\% \\ BR(H \rightarrow \gamma\gamma) &= 0.24\% \end{aligned}$$

$$\begin{aligned} BR(H \rightarrow bb) &= 56.9\% \\ BR(H \rightarrow \tau\tau) &= 6.2\% \\ BR(H \rightarrow \mu\mu) &= 0.022\% \end{aligned}$$

→ at 125 GeV: only ~11% of decays not observable (gg, cc)

\*) LHC Higgs cross-section working group

# Status of Higgs boson physics at the LHC




Expected number of decays, before selection  
cuts, in the data,  $m_H = 125$  GeV:

- ~ 950  $H \rightarrow \gamma\gamma$
- ~ 60  $H \rightarrow ZZ^* \rightarrow 4\ell$
- ~ 9000  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$




# Result of the Searches for $H \rightarrow \gamma\gamma$

Phys. Rev. D90 (2014) 112015



EPJ C74 (2014) 3076




- Background interpolation in the region of the excess (obtained from sidebands)
- Reducible  $\gamma$ -jet and jet-jet background at the level of 25%
- High signal significance in both experiments: ATLAS:  $5.2\sigma$  ( $4.6\sigma$  expected)  
CMS:  $5.7\sigma$  ( $5.2\sigma$  expected)
- Establishes the discovery in this channel alone




# Result of the Searches for $H \rightarrow \gamma\gamma$

Phys. Rev. D90 (2014) 112015



EPJ C74 (2014) 3076




Measured signal strengths:  $\mu = \sigma_{\text{obs}} / \sigma_{\text{SM}}$

ATLAS:  $\mu = 1.17 \pm 0.27$


CMS:  $\mu = 1.14 \pm 0.26$



# Categorisation of $H \rightarrow \gamma\gamma$ candidate events




Categorisation: to increase overall sensitivity and sensitivity to different production modes (VBF, VH)



- VH enriched: one-lepton,  $E_T^{\text{miss}}$ , low-mass di-jets
- VBF enriched (tag-jet configuration,  $\Delta\eta$ ,  $m_{jj}$ )
- gluon fusion: exploit different mass resolution for different detector regions,  $\gamma\gamma$  conversion status and  $p_{Tt}$


# $H \rightarrow \gamma\gamma$ VBF candidate event



Run Number: 204769, Event Number: 24947130

Date: 2012-06-10 08:17:12 UTC

# $H \rightarrow \gamma\gamma$ VBF candidate event



$$E_T(\gamma_1) = 80.1 \text{ GeV}, \eta = 1.01$$


$$E_T(\gamma_2) = 36.2 \text{ GeV}, \eta = 0.17$$

$$m_{\gamma\gamma} = 126.9 \text{ GeV}$$

$$E_T(\text{jet}_1) = 121.6 \text{ GeV}, \eta = -2.90$$

$$E_T(\text{jet}_2) = 82.8 \text{ GeV}, \eta = 2.72$$

$$m_{jj} = 1.67 \text{ TeV}$$





**ATLAS**  
**EXPERIMENT**

Run Number: 204769, Event Number: 24947130

Date: 2012-06-10 08:17:12 UTC


# $\gamma\gamma$ signal strengths for various production modes



Fit results for individual production processes are consistent with the Standard Model expectations

# $H \rightarrow ZZ \rightarrow e^+e^- \mu^+\mu^-$ candidate event

 **ATLAS**  
EXPERIMENT  
<http://atlas.ch>




Run: 182796  
Event: 74566644  
2011-05-30 07:54:29 CEST


# Reconstructed mass spectra from $4\ell$ decays



Phys. Rev. D91 (2014) 012006



Phys. Rev. D89 (2014) 092007



Measured signal strengths:


ATLAS:  $\mu = 1.44^{+0.40}_{-0.33}$

CMS:  $\mu = 0.93^{+0.29}_{-0.23}$


Significance in each experiment  $> 6\sigma$

# $H \rightarrow WW^* \rightarrow \ell\nu \ell\nu$ signal

arXiv:1412.2641



JHEP 01 (2014) 096



Measured signal strengths:

ATLAS:  $\mu = 1.09^{+0.23}_{-0.21}$


CMS:  $\mu = 0.72^{+0.20}_{-0.18}$

- Very significant excesses visible in the “transverse mass” (ATLAS:  $6.1\sigma$ ) and  $m_{\ell\ell}$  distributions (CMS:  $4.5\sigma$ )

# $H \rightarrow WW^* \rightarrow l\nu l\nu$ signal



arXiv:1412.2641




Measured signal strengths: ATLAS

Gluon fusion (ggF):  $\mu = 1.02^{+0.29}_{-0.26}$


VBF:  $\mu = 1.27^{+0.53}_{-0.45}$

# Differential cross-section measurements

JHEP 09 (2014) 112




Phys. Lett B738 (2014) 234




- First fiducial, differential cross-section measurements in bosonic channels
- Good agreement within present experimental and theoretical uncertainties, (... except normalization?)
- Large future potential: probe Higgs boson kinematics, jet activity, VBF contributions, spin-CP nature, ...

# Couplings to quarks and leptons ?




- Search for  $H \rightarrow \tau\tau$  and  $H \rightarrow bb$  decays;
- Challenging signatures due to jets ( $bb$  decays) or significant fraction of hadronic tau decays
- Vector boson fusion mode essential for  $H \rightarrow \tau\tau$  decays





- Associated production  $WH$ ,  $ZH$  modes have to be used for  $H \rightarrow bb$  decays



- Exploitation of multivariate analyses



# Evidence for $H \rightarrow \tau\tau$ decays



$m_{\tau\tau}$  distribution, events weighted by  $\ln(1+S/B)$


Measured signal strengths:

ATLAS:  $\mu = 1.43^{+0.43}_{-0.37}$   $(4.5\sigma)$

CMS:  $\mu = 0.78 \pm 0.27$   $(3.2\sigma)$

*One of the most important LHC results in 2014*

# Results on the search for $H \rightarrow bb$ decays



Reconstructed  $m_{bb}$  signals (after subtraction of major, non-resonant backgrounds)

- Reference signal from WZ, and ZZ with  $Z \rightarrow bb$  seen
- Positive, but non-conclusive Higgs boson signal contribution observed

## Signal strengths:



ATLAS:  $\mu = 0.50 \pm 0.36$

CMS:  $\mu = 1.0 \pm 0.5$

# Results on the search for $H \rightarrow \mu\mu$ decays



Phys. Lett. B738 (2014) 68



$m_H = 125$  GeV:

ATLAS 95% CL:  $7.0\sigma_{SM}$  (7.2 expected, no Higgs)

[Phys. Lett. B738 (2014) 68]

CMS 95% CL:  $7.4\sigma_{SM}$  (6.5 expected, no Higgs)

[Phys. Lett. B744 (2015) 184]

$\rightarrow BR(H \rightarrow \mu\mu) < \sim 1.5 \cdot 10^{-3}$

Significantly smaller than  $BR(H \rightarrow \tau\tau)$




$\rightarrow$  no evidence for flavour-universal coupling

# Lepton-flavour violating $H \rightarrow \tau\mu$ decays?



arXiv:1502.07400


Search for  $\tau \rightarrow \text{had } \nu$  or  $\tau \rightarrow e\nu\nu$   
accompanied by 0, 1 or 2 jets




- CMS analysis: a slight excess of events observed ( $2.4\sigma$ )
- Consistent with a signal at 125 GeV with a  $H \rightarrow \tau\mu$  branching ratio of  $(0.84^{+0.39}_{-0.37})\%$
- ATLAS results on this search are eagerly awaited, expected soon

# Search for invisible Higgs boson decays

- Some extensions of the Standard Model allow a Higgs boson to decay to stable or long-lived particles




- Search for excess in ZH associated production and VBF production



**VBF**

**ZH**

**PRL 112 (2014) 201802**



Assuming the ZH and VBF production rates for  $m_H = 125 \text{ GeV}$ :

ATLAS: 95% CL on  $\text{BR}(H \rightarrow \text{inv.}) < 0.75$  (from ZH production)


95% CL on  $\text{BR}(H \rightarrow \text{inv.}) < 0.29$  (from VBF production) [ATLAS-CONF-2015-004]

CMS: 95% CL on  $\text{BR}(H \rightarrow \text{inv.}) < 0.58$  (from ZH + VBF combination)


# Interpretation in Higgs-portal models

-Stable dark matter particles with couplings to the Higgs boson-

- For  $m_\chi < m_H/2$ , limits on invisible branching ratios can be translated to the spin-independent DM-nucleon elastic cross section for scalar, vector and fermionic DM particles
- Higgs-nucleon coupling, model dependent: assume  $0.33^{+0.30}_{-0.07}$  (lattice calculations)
- Within this model, interesting limits for low  $m_\chi$  masses



Parallel Session Talks: James Beacham






# *Profile of the New Particle*

## *Is it the Standard Model Higgs Boson?*

- Mass (“input parameter”)
- Width
- Spin,  $J^{CP}$  quantum number
- Production rates

### Couplings to bosons and fermions







### Parallel Session Talks:

- Marcello Fanti  
(couplings / properties)
- Daniele Zanzi (ttH)

# Higgs boson mass

- The two high resolution channels  $H \rightarrow ZZ^* \rightarrow 4\ell$  and  $H \rightarrow \gamma\gamma$  are best suited (reconstructed mass peak, good mass resolution)
- Good control of the lepton and photon energy scales, calibration via  $Z \rightarrow \ell\ell$  and  $J/\psi$  and  $\Upsilon$  signals, improved understanding of lepton and photon reconstruction



ATLAS:  $m_H = 125.36 \pm 0.37 \text{ (stat)} \pm 0.18 \text{ (syst)} \text{ GeV}$

CMS:  $m_H = 125.02^{+0.26}_{-0.27} \text{ (stat)}^{+0.14}_{-0.15} \text{ (syst)} \text{ GeV}$




# Higgs boson mass (cont.)

-First ATLAS and CMS combination of Higgs boson results-



PRL 114 (2015) 191803

## Individual and combined results:




## ATLAS + CMS:

$$m_H = 125.09 \pm 0.21 \text{ (stat)} \pm 0.11 \text{ (syst)} \text{ GeV}$$

Precision of 0.2%


## Uncertainties:



- Statistical uncertainty still dominant
- Major systematic uncertainties: Lepton and photon energy scales and resolutions
- Theoretical uncertainties small (correlated),  $\gamma\gamma$  interference effects neglected

# Higgs boson width

- The Standard Model Higgs boson width is expected to be small:  $\Gamma_H \sim 4 \text{ MeV}$
- Experimental mass resolution in  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4\ell$  channel  $\sim 1 - 2 \text{ GeV}$   
→ only upper limits can be extracted from the observed mass peaks



EPJ C75 (2015) 5, 212






Results: 95% CL limits

$\Gamma_H < 1.7 \text{ GeV}$  (2.3 expected)

# Indirect constraint on the Higgs boson width from “off-shell cross sections”

- Different sensitivity of on-shell and off-shell cross sections on the Higgs boson width
- However, model dependent: assumes that on-shell and off-shell couplings are the same
- Dependence on K-factors for signal and backgrounds ( $gg \rightarrow VV$ )



**Results:** 95% CL limits



**CMS:**  $\Gamma_H / \Gamma_{SM} < 5.4$  ( $= 22 \text{ MeV}$ )

**ATLAS:**  $\Gamma_H / \Gamma_{SM} < 5.5$  ( $R_B^{H^*} = 1$ )

# Spin and CP

- Standard Model Higgs boson:  $J^P = 0^+$   
→ strategy is to falsify other hypotheses  
( $0^-$ ,  $1^-$ ,  $1^+$ ,  $2^-$ ,  $2^+$ )
- Angular distributions of final state particles show sensitivity to spin

In particular:  $H \rightarrow ZZ^* \rightarrow 4\ell$  decays  
(in addition:  $H \rightarrow WW^* \rightarrow \ell\nu \ell\nu$ )




- Data strongly favour the spin-0 hypothesis of the Standard Model
- Many alternatives can be excluded with confidence levels  $> 99\%$


# Result on different $J^{CP}$ hypothesis tests



Phys. Lett. B726 (2013) 120





arXiv:1411.3441



- In both experiments, data are consistent with  $J^P = 0^+$  hypothesis, many alternative models are excluded with high significance

# Signal strength in individual decay modes

-normalised to the expectations for the Standard Model Higgs boson-



Signal strengths:

ATLAS:  $\mu = 1.18^{+0.15}_{-0.14}$



CMS:  $\mu = 1.00 \pm 0.14$

- Data are consistent with the hypothesis of the Standard Model Higgs boson
- If ATLAS and CMS combined: clear evidence for coupling to fermions

# Higgs boson couplings



- Production and decay involve several couplings

Production:



Decays: e.g  $H \rightarrow \gamma\gamma$  (best example)

(Decay widths depends on W and top-coupling, destructive interference)



- Benchmarks defined by LHC cross section working group (leading-order tree-level framework):

- Signals observed originate from a single resonance;  
(mass assumed here is 125.36GeV (ATLAS) and 125.02 GeV (CMS))
- Narrow width approximation: → rates for given channels can be decomposed as:

$$\sigma \cdot B(i \rightarrow H \rightarrow f) = \frac{\sigma_i \cdot \Gamma_f}{\Gamma_H}$$

$i, f$  = initial, final state  
 $\Gamma_f, \Gamma_H$  = partial, total width




- Modifications to coupling strength are considered (coupling scale factors  $\kappa$ ), tensor structure of Lagrangian assumed as in Standard Model

# Couplings to fermions and bosons

- Assume only one scale factor for fermion and vector couplings:

$$\kappa_V = \kappa_W = \kappa_Z \quad \text{and} \quad \kappa_F = \kappa_t = \kappa_b = \kappa_\tau$$

- Assume that  $H \rightarrow \gamma\gamma$  and  $gg \rightarrow H$  loops and the total Higgs boson width depend only on  $\kappa_V$  and  $\kappa_F$  (no contributions from physics beyond the Standard Model)
- Sensitivity to relative sign between  $\kappa_F$  and  $\kappa_V$  only from interference term in  $H \rightarrow \gamma\gamma$  decays (assume  $\kappa_V > 0$ )




# Higgs boson couplings

- Fit all coupling scale factors for relevant particles ( $W, Z, t, b, \tau, \mu$ ) independently;
- Loop factors expressed in terms of these scale factors, assume SM particle content




$\lambda$  = Yukawa coupling for fermions  
 $\sqrt{g}/2v$  = couplings for  $W/Z$  bosons

EPJ C75 (2015) 5, 212




ATLAS-CONF-2015-007




For the first time, non-universal, mass-dependent couplings observed

# Ratios of Higgs boson couplings (model independent)

- In the most general model, only ratios of couplings can be measured independently on any assumptions on the total width (allowing also deviations in vertex loop coupling strength)



$\lambda_{WZ}$ : test of custodial symmetry  
 $\lambda_{\gamma Z}$ : sensitive to new charged particles in  $H \rightarrow \gamma\gamma$  loop w.r.t.  $H \rightarrow ZZ$  decays  
 $\lambda_{tg}$ : sensitive to new coloured particles contributing to  $gg \rightarrow H$  production w.r.t.  $t\bar{t}H$  production



Good consistency with the Standard Model Higgs boson hypothesis

# *Additional Higgs bosons?*

*Composite  
Higgs bosons*

*More Higgs bosons*

*SUSY Higgs*

*No Higgs at the LHC*

*MSSM Higgs bosons*

*Dark Higgs*

*Heidi Higgs*




Parallel Session Talks:

- Matthias Mozer
- Nicolaos Raptopotis
- Glauber Dorsch

# Search for Additional Higgs Bosons

## -a few examples-



(i) Results of an ATLAS search on additional resonances X decaying into  $\gamma\gamma$



Observed and expected 95% CL limits on the fiducial cross section times branching ration  $\text{BR}(X \rightarrow \gamma\gamma)$  as a function of mass

(note: 125 GeV signal was treated as “background” and contribution was subtracted)


(ii) Results of a CMS search on additional SM-like Higgs bosons decaying into ZZ and WW



Observed and expected 95% CL limits on the cross section normalised to the SM value for individual channels and their combination


### (iii) Search for charged and heavy neutral MSSM Higgs bosons

Search for  $H^\pm \rightarrow \tau\nu$  decays via  $t\bar{t}$  production or  $tH^\pm$  associated production



95% CL exclusion limits on branching ratios or cross sections times branching ratio


Parallel Session Talks: Nicolaos Rompotis



Expected and observed exclusion limits at 95% CL in the  $(m_A - \tan\beta)$  parameter plane for the MSSM  $m_h^{\text{mod+}}$  benchmark scenario

# What next?

The LHC has started operation at  $\sqrt{s} = 13$  TeV




A proton-proton collision at 13 TeV, recorded by the ATLAS experiment on 21. May 2015

# A new energy range will be explored !!


Major physics topics:

- (i) Extend the searches for New Physics
- (ii) Precise measurements of the Higgs boson profile
- (iii) Additional Higgs bosons?
- (iv) Scattering of vector bosons
- (v) Precision measurements ( $m_W$ ,  $m_{top}$ , Higgs couplings)





# First evidence for electroweak $W^\pm W^\pm jj$ production (Run 1)


- Higgs boson needed in the SM to regularise VV scattering at high energies;
- Key experimental process:  $W^\pm W^\pm$  scattering



Electroweak production



Strong production



VBS enhancement by cutting on mass ( $m_{jj}$ ) and rapidity separation  $\Delta y_{jj}$



ATLAS:  $3.6\sigma$  for electroweak production

CMS:  $2.0\sigma$

PRL 114 (2015) 051801

(expected: in both experiments about  $3\sigma$ )

# Conclusions

- The analyses of the complete LHC Run 1 dataset by the ATLAS and CMS experiments have consolidated the milestone discovery announced in July 2012
  - Properties of the particle ( $J^{CP}$ , couplings) are in very good agreement with those expected for the Standard Model Higgs boson
- The experiments have moved from the discovery to the measurement phase;
- Many measurements still statistically limited  
→ significant improvements expected in Run 2 and beyond
  - → Higgs particle might be the portal to new physics
  - Exciting times ahead of us, with new, unexplored energy regime in reach

